



Transverse mass distributions of strange particles produced in Pb-Pb collisions at 158 A GeV/c

G. E. Bruno^b for the NA57 Collaboration:

F Antinori^l, P Bacon^e, A Badalà^g, R Barbera^g, A Belogianni^a, A Bhasin^e, I J Bloodworth^e, M Bombari^j, G E Bruno^b, S A Bull^e, R Caliendo^b, M Campbell^h, W Carena^h, N Carrer^h, R F Clarke^e, A Dainese^l, A P de Haas^s, P C de Rijke^s, D Di Bari^b, S Di Liberto^o, R Divia^h, D Elia^b, D Evans^e, K Fanebust^d, G A Feofilov^q, R A Fini^b, P Ganoti^a, B Ghidini^b, G Grella^p, H Helstrup^d, A K Holme^k, A Jacholkowski^b, G T Jones^e, P Jovanovic^e, A Juskoⁱ, R Kamermans^s, J B Kinson^e, K Knudson^h, A A Kolozhvari^q, V Kondratiev^q, I Králikⁱ, A Kravčáková^j, P Kuijer^s, V Lenti^b, R Lietava^f, G Løvholden^k, V Manzari^b, G Martinská^j, M A Mazzoni^o, F Meddi^o, A Michalon^r, M Morando^l, F Navach^b, P I Norman^e, A Palmeri^g, G S Pappalardo^g, B Pastirčákⁱ, J Pišúf^f, N Pisutova^f, E Quercigh^l, F Riggi^g, D Röhrich^c, G Romano^p, K Šafařík^h, L Šándorⁱ, E Schillings^s, G Segato^l, M Senè^m, R Senè^m, W Snoeys^h, F Soramel^l, M Spyropoulou-Stassinaki^a, P Starobaⁿ, T A Toulina^q, R Turrisi^l, T S Tveter^k, J Urbán^j, F F Valiev^q, A van den Brink^s, P van de Ven^s, P Vande Vyvre^h, N van Eijndhoven^s, J van Hunen^h, A Vascotto^h, T Vik^k, O Villalobos Baillie^e, L Vinogradov^q, T Virgili^p, M F Votruba^e, J Vrláková^j and P Závadaⁿ

^a Physics Department, University of Athens, Athens, Greece

^b Dipartimento IA di Fisica dell'Università e del Politecnico di Bari and INFN, Bari, Italy

^c Fysisk Institutt, Universitetet i Bergen, Bergen, Norway

^d Høgskolen i Bergen, Bergen, Norway

^e University of Birmingham, Birmingham, UK

^f Comenius University, Bratislava, Slovakia

^g University of Catania and INFN, Catania, Italy

^h CERN, European Laboratory for Particle Physics, Geneva, Switzerland

ⁱ Institute of Experimental Physics, Slovak Academy of Science, Košice, Slovakia

^j P.J. Šafařík University, Košice, Slovakia

^k Fysisk Institutt, Universitetet i Oslo, Oslo, Norway

^l University of Padua and INFN, Padua, Italy

^m Collège de France, Paris, France

ⁿ Institute of Physics, Prague, Czech Republic

^o University “La Sapienza” and INFN, Rome, Italy

^p Dipartimento di Scienze Fisiche “E.R. Caianiello” dell'Università and INFN, Salerno, Italy

^q State University of St. Petersburg, St. Petersburg, Russia

^r Institut de Recherches Subatomique, IN2P3/ULP, Strasbourg, France

^s Utrecht University and NIKHEF, Utrecht, The Netherlands

Experiment NA57 has collected high statistics, high purity samples of K_S^0 , Λ , Ξ and Ω produced in Pb-Pb collisions at 158 A GeV/c. In this paper we present a study of the transverse mass spectra of these particles for a sample of events corresponding to about the most central 55% of the inelastic Pb-Pb cross section. We analyse the transverse mass distributions in the framework of the blast-wave model for the full sample under consideration and, for the first time at the SPS, as a function of the event centrality.

1 Introduction

The NA57 experiment has been designed to study the production of strange baryons and antibaryons in heavy ion collisions. The experiment has extended the WA97 [1] measurements to a wider centrality range and to a lower beam momentum. NA57 results on hyperon yields [2] at 158 A GeV/c confirm the enhancement pattern observed by the WA97 experiment, with an enhancement factor increasing with increasing strangeness. In this paper we shall concentrate on the analysis of the

transverse mass ($m_T = \sqrt{p_T^2 + m^2}$) spectra for the Λ , Ξ^- , Ω^- hyperons, their antihyperons and K_S^0 measured in Pb-Pb collisions at 158 A GeV/c.

2 The NA57 experiment

The layout of the NA57 experiment has been described in detail elsewhere [3]. A telescope of compactly packed silicon pixel detectors is used as main tracking device. Additional pixel and double-sided silicon microstrip detectors

are used to improve the momentum resolution for high momentum tracks. The telescope inclination angle with respect to the beam line and the distance from the target are set to accept particles produced in about half a unit of rapidity around central rapidity and medium transverse momentum.

An array of scintillation counters provides a fast signal to trigger on the centrality of the collisions. A more precise centrality measurement is provided by two stations of silicon strip detectors. All detectors are located in the 1.4 T field of the Goliath magnet. The triggered fraction of the total inelastic cross section is about 55%.

The K_S^0 mesons, the Λ , Ξ^- and Ω^- hyperons and their antiparticles are identified by reconstructing their weak decays into final states containing only charged particles. Decays are required to take place in a fiducial volume located between a fixed distance from the target and the first plane of the telescope. The selection procedure allows for extraction of hyperon and K_S^0 signals with negligible background [4].

Events have been divided into five centrality classes (0,1,2,3,4), class 0 being the most peripheral and class 4 the most central, according to the charge particle multiplicity measured in the pseudorapidity intervals $2 < \eta < 3$ and $3 < \eta < 4$. The centrality of the collision is expressed as number of wounded nucleons computed from the measured trigger cross sections using the Glauber model. The centrality determination is described in detail in ref. [5].

3 Data sets and analysis

The results presented in this paper are based on the analysis of a data sample consisting of 460 M events of Pb-Pb collisions collected in the years 1998 and 2000.

For each particle species we define the fiducial acceptance window using a Monte Carlo simulation of the apparatus, in order to exclude the border regions.

All data are corrected for geometrical acceptance and for detector and reconstruction inefficiencies on a particle-by-particle basis, using a Monte Carlo technique where simulated particles are embedded into real events (the procedure has been described in detail in [6]).

The correction method is CPU intensive; therefore, while all the reconstructed Ξ s and Ω s have been individually weighted, for the much more abundant K_S^0 , Λ and $\bar{\Lambda}$ samples we only weighted a small fraction of the total sample in order to reach a statistical accuracy better than the limits imposed by the systematic error.

4 Inverse slopes of the transverse mass spectra

The double-differential (y, m_T) distributions for each particle have been parametrized using the expression

$$\frac{d^2N}{m_T dm_T dy} = f(y) \exp\left(-\frac{m_T}{T_{app}}\right) \quad (1)$$

assuming the rapidity distribution to be flat ($f(y) = \text{const}$) within our acceptance region. The inverse slope parameter T_{app} (“apparent temperature”) has been extracted by means of a maximum likelihood fit of Eq. 1 to the data. As discussed in detail in the next section, this apparent temperature can be interpreted as due to the thermal motion coupled with a collective transverse flow of the fireball components assumed to be in thermal equilibrium.

Transverse mass distributions $1/m_T dN/dm_T$ for strange particles measured in the centrality range accessible to the experiment are shown in figure 1. The shapes of all spectra




Figure 1. Transverse mass spectra of strange particles from about the 55% most central Pb-Pb cross section at 158 A GeV/c. The superimposed exponential functions have inverse slopes equal to the T_{app} values obtained from the maximum likelihood fits.

are well described by single exponential functions. In the next section, we shall attempt a global fit with the blast-wave model.

The inverse slope parameters T_{app} of the transverse mass spectra are given in table 1. They are in agreement within the errors with those measured over a smaller centrality range (about 40% of the Pb-Pb inelastic cross section) by the WA97 experiment [7].

Table 1. Inverse slopes parameter T_{app} (MeV) of the strange particles in the full centrality range (0–4). The first error is statistical, the second one systematic.

	Inverse Slope (MeV)
K_S^0	$237 \pm 4 \pm 24$
Λ	$289 \pm 7 \pm 29$
$\bar{\Lambda}$	$287 \pm 6 \pm 29$
Ξ^-	$297 \pm 5 \pm 30$
$\bar{\Xi}^+$	$316 \pm 11 \pm 30$
Ω^-	$264 \pm 19 \pm 27$
$\bar{\Omega}^+$	$284 \pm 28 \pm 27$

5 Blast-wave description of the spectra

We consider in this section the statistical hadronization model of ref. [8] to describe the strange particle spectra discussed above. The model assumes that particles decouple from a system in local thermal equilibrium with temperature T , which expands both longitudinally and in the transverse direction; the quantum statistical distributions are approximated by the Boltzmann distribution. The longitudinal expansion is boost-invariant, the transverse one is defined in terms of a transverse velocity field.

The blast-wave model predicts a double differential cross section for a given particle j of the form:

$$\frac{d^2 N_j}{m_T dm_T dy} = \int_0^{R_G} \mathcal{A}_j m_T K_1 \left(\frac{m_T \cosh \rho}{T} \right) I_0 \left(\frac{p_T \sinh \rho}{T} \right) r dr \quad (2)$$

where $\rho(r) = \tanh^{-1} \beta_\perp(r)$, K_1 and I_0 are two modified Bessel functions and R_G is the transverse geometric (gaussian) radius of the source.

In a preliminary study [9], we considered a simplified model where the transverse flow velocity was assumed to be constant all along its profile: $\beta_\perp(r) = \langle \beta_\perp \rangle$. A more detailed analysis using different transverse velocity profiles has now been performed. We have also studied the stability of the fits (the parameters T and $\langle \beta_\perp \rangle$ are strongly anti-correlated), as well as their sensitivity to individual particle spectra. A complete description of the transverse mass spectrum analysis at 160 A GeV/c will be presented in a forthcoming publication [10]. In this new study, the transverse velocity field $\beta_\perp(r)$ has been parametrized according to power laws:

$$\beta_\perp(r) = \beta_S \left[\frac{r}{R_G} \right]^n \quad r \leq R_G \quad (3)$$

where the exponent n was assumed to be $n = 0, 1/2, 1$ or 2 . Assuming a uniform particle density, the average transverse velocity is calculated according to $\langle \beta_\perp \rangle = \frac{2}{2+n} \beta_S$. As shown in fig. 2, the simultaneous best fit of eq. 2 ($n = 1$ profile case) to the data points of all the measured

strange particle spectra successfully describes the data with $\chi^2/ndf = 37.2/48$, yielding the following values for the two basic quantities T and $\langle \beta_\perp \rangle$:

$$T = 144 \pm 7(\text{stat}) \pm 14(\text{syst}) \text{ MeV}$$

$$\langle \beta_\perp \rangle = 0.381 \pm 0.013(\text{stat}) \pm 0.012(\text{syst}).$$

The quadratic profile is disfavoured by the data, yielding the largest value of χ^2 . The other two profiles ($n = 0, n = 1/2$) yield similar values of the freeze-out temperatures and of the average transverse flow velocities, as the $n = 1$ case, with good values of χ^2/ndf . For the study of the centrality dependence of the freeze-out parameters presented below, the linear profile has been used.




Figure 2. Transverse mass spectra fitted with collective flow model for the linear ($n = 1$) transverse velocity profile. The quoted values are valid for the full NA57 centrality sample.

The particles have been divided into two groups — those which share quarks in common with the nucleons and those which do not¹ — and the fit procedure has been repeated separately for the two groups. The results of such fits are summarized in table 2. They suggest similar freeze-out conditions for the two groups. Since the interaction cross-sections for the particles of the two groups are quite different, this finding would suggest a similar production mechanism and final state interactions of limited importance (e.g. a rapid thermal freeze-out). A similar conclusion concerning the evolution of the system was reached in the study of the HBT correlation functions of negative pions at high p_T in the WA97 experiment [11].

¹It is known that the particles of the two groups may exhibit different production features, e.g. in the rapidity spectra.

Table 2. Thermal freeze-out temperature T and average transverse flow velocity $\langle \beta_{\perp} \rangle$ in the full centrality range. The first error is statistical, the second one systematic.

	T (MeV)	$\langle \beta_{\perp} \rangle$	χ^2/ndf
K_S^0, Λ, Ξ^-	$146 \pm 8 \pm 14$	$0.38 \pm 0.01 \pm 0.01$	18.1/23
$\bar{\Lambda}, \bar{\Xi}^+, \Omega^-, \bar{\Omega}^+$	$130 \pm 28 \pm 14$	$0.40 \pm 0.03 \pm 0.01$	18.5/23

In order to see whether some species deviate from the common temperature and transverse flow velocity scenario determined above, we have attempted to fit the spectra separately. For statistical reasons, Ω^- and $\bar{\Omega}^+$ spectra have been combined. For all particle species, T and $\langle \beta_{\perp} \rangle$ do not deviate more than two standard deviations from the common freeze-out scenario.

We have also performed the global fit to the spectra for each of the five NA57 centrality classes. The results are summarized in table 3.

Table 3. Thermal freeze-out temperature (T) and average transverse flow velocity ($\langle \beta_{\perp} \rangle$) in five centrality classes, assuming a linear transverse profile.

	T (MeV)	$\langle \beta_{\perp} \rangle$	χ^2/ndf
0	139 ± 25	0.35 ± 0.04	33.2/35
1	151 ± 15	0.35 ± 0.03	42.3/42
2	146 ± 17	0.37 ± 0.03	41.3/43
3	135 ± 14	0.41 ± 0.02	31.8/41
4	118 ± 13	0.45 ± 0.02	52.7/43

In figure 3 we show the 1σ confidence level contours in the $\langle \beta_{\perp} \rangle$ - T parameter space as obtained from the blast-wave fit with the $n = 1$ profile case. Apart from the most peripheral interactions, the centrality dependence of the two parameters shows opposite trends, the temperature is decreasing and the average flow velocity is increasing with increasing centrality.

6 Conclusions and outlook

We have analysed the transverse mass spectra of high statistics, high purity samples of K_S^0 , Λ , Ξ and Ω particles produced in Pb-Pb collisions at 158 A GeV/c over a wide range of collision centrality (about the most central 55% of the Pb-Pb interaction cross section).

For each hyperon species, the m_T slope is found to be in good agreement with that of the corresponding anti-hyperon.

The analysis in the framework of a blast-wave model using a linear velocity profile suggests that after a central collision the system expands explosively; the system then freezes-out when the temperature is of the order of 120




Figure 3. The 1σ confidence level contours from fits to individual centrality classes..

MeV with an average transverse flow velocity of about one half of the speed of light. Particles with and without valence quarks in common with the nucleon appear to have similar behaviour. With increasing collision centrality, the transverse flow velocity increases and the final temperature decreases.

References

1. E. Andersen *et al.* [WA97 Collaboration], Phys. Lett. B **449** (1999) 401
2. L. Sándor *et al.* [NA57 Collaboration], Proceeding of the 7th International Conference on Strange Quarks in Matter, to be published in J. Phys. G: Nucl. Part. Phys. (2003)
3. V. Manzari *et al.* [NA57 Collaboration], J. Phys. G: Nucl. Part. Phys. **25** (1999) 473
4. V. Manzari *et al.* [NA57 Collaboration], Nucl. Phys. A **661** 761c
5. N. Carrer *et al.* [NA57 Collaboration], J. Phys. G: Nucl. Part. Phys. **27** (2001) 391
6. K. Fanebust *et al.* [NA57 Collaboration], J. Phys. G: Nucl. Part. Phys. **28** (2002) 160
7. F. Antinori *et al.* [WA97 Collaboration], Eur. Phys. J. C **14** (2000) 633
8. E. Schnedermann, J. Sollfrank and U. Heinz, Phys. Rev. C **48** (1993) 2462
9. G. E. Bruno *et al.* [NA57 Collaboration], Proceeding of the XXXVIII Rencontres de Moriond (2003) Preprint nucl-ex/0305033
10. F. Antinori *et al.* [NA57 Collaboration], being submitted to J. Phys. G: Nucl. Part. Phys.
11. F. Antinori *et al.* [WA97 Collaboration], J. Part. Phys. G: Nucl. Part. Phys. **27** (2001) 2325.